



**(43) International Publication Date**  
**19 August 2004 (19.08.2004)**

**PCT**

**(10) International Publication Number**  
**WO 2004/069332 A1**


- (51) International Patent Classification<sup>7</sup>: A61N 1/36
- (21) International Application Number: PCT/US2004/002849
- (22) International Filing Date: 30 January 2004 (30.01.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- |            |                              |    |
|------------|------------------------------|----|
| 10/358,093 | 3 February 2003 (03.02.2003) | US |
| 10/752,940 | 6 January 2004 (06.01.2004)  | US |
- (71) Applicant (*for all designated States except US*): EN-TEROMEDICS INC. [US/US]; 2800 Patton Road, St. Paul, MN 55113 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): KNUDSON, Mark,

**B. [US/US];** 1309 West Royal Oaks Drive, Shoreview, MN 55126 (US). **CONRAD, Timothy, R. [US/US];** 12557 Riverview Road, Eden Prairie, MN 55347 (US). **EVNIN, Luke, B. [US/US];** 170 Belgrave Avenue, San Francisco, CA 94117 (US). **WILSON, Richard, R. [US/US];** 1276 Nursery Hill Lane, Arden Hills, MN 55112 (US). **TWEDEN, Katherine, S. [US/US];** 1175 Ashley Lane, Mahtomedi, MN 55115 (US).

- (74) **Agent:** BRUESS, Steven, C.; Merchant & Gould P.C.,  
P.O. Box 2903, Minneapolis, MN 55402-0903 (US).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,

[Continued on next page]

- (54) Title: INTRALUMINAL ELECTRODE



**(57) Abstract:** At least one of a plurality of disorders of a patient associated with vagal activity innervating at least one of a plurality of organs of the patient at an innervation site are treated by positioning a neurostimulator carrier within a body lumen of the patient. An electrode disposed on the carrier is positioned at a mucosal layer of the lumen. An electrical signal is applied to the electrode to modulate vagal activity by an amount selected to treat the disorder. The signal may be a blocking or a stimulation signal.

**WO 2004/069332 A1**

AP3



TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- *with international search report*

- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## INTRALUMINAL ELECTRODE

This application is being filed on 30 January 2004, as a PCT International Patent application in the name of EnteroMedics Inc., a U.S. national corporation, applicant for all countries except the US, and Mark B. Knudson, Timothy R. Conrad, Luke B. Evnin, Richard R. Wilson, and Katherine S. Tweden, all US citizens, applicants for the designation of the US only.

### **BACKGROUND OF THE INVENTION**

#### 10      **1. Field of the Invention**

This invention pertains to treatments of disorders associated, at least in part, with neural activity. These may include, without limitation, gastrointestinal, pancreo-biliary, cardio-respiratory and central nervous system disorders (including neurological and psychiatric, psychological and panic disorders). More particularly, 15 this invention pertains to treatment of such disorders through management of neural impulse stimulation and blocking.

#### **2. Description of the Prior Art**

##### **A. Functional Gastrointestinal Disorders (FGIDs)**

20      Functional Gastrointestinal Disorders (FGIDs) are a diagnostic grouping having diagnostic criteria based on symptomatology, because the pathophysiology of these diseases is multifactorial with some pathophysiologic mechanisms in common. FGIDs are thought to be due to altered autonomic nervous system balance and to be pathophysiological combinations of: (1) abnormal GI motility; (2) visceral 25 hypersensitivity; and, (3) brain-gut interactions. Tougas, "The Autonomic Nervous System in Functional Bowel Disorders", Gut, Vol. 47 (Suppl IV), pp. iv78-iv80 (2000) and Drossman, "Rome II: A Multinational Consensus Document on Gastrointestinal Disorders – The Functional Gastrointestinal Disorders and the Rome II Process", Gut, Vol. 45 (Suppl II):II1-II5 (1999). The FGIDs of interest to 30 the present invention are functional dyspepsia (dysmotility-like) and irritable bowel syndrome (IBS).

##### **1. Functional Dyspepsia (Dysmotility-Like)**

35      Functional dyspepsia (dysmotility-like), is diagnosed when a patient's symptoms, in the absence of other organic disease likely to explain the symptoms, include persistent or recurrent pain or discomfort centered in the upper abdomen that may be accompanied by upper abdominal fullness, early satiety, bloating or nausea. Talley et al., "Rome II: A Multinational Consensus Document on Gastrointestinal

Disorders – Functional Gastrointestinal Disorders” Gut, Vol. 45 (Suppl II), pp. II37-II42 (1999).

A spectrum of dysmotilities has been documented in patients with functional dyspepsia. These include delayed gastric emptying of solids and liquids, reduced  
 5 vagal tone, gastric dysrhythmias and impaired gastric accommodation. Furthermore, some studies have found good correlation between symptoms and indices of dysmotility, while others have not. Stanghellini V, et al., “Delayed Gastric Emptying of Solids in Patients with Functional Dyspepsia”, Gastroenterol, (1996) 110:1036-1042. Undeland KA, et al., “Wide Gastric Antrum and Low Vagal Tone  
 10 in Patients with Diabetes Mellitus Type 1 Compared to Patients with Functional Dyspepsia and Healthy Individuals”, Dig Dis Sci, (1996) 41:9-16. Tack J, et al., “Role of Impaired Gastric Accommodation to a Meal in Functional Dyspepsia”, Gastroenterol, (1998) 115:1346-1352. Wilmer A, et al., “Ambulatory Gastrojejunal Manometry in Severe Motility-like Dyspepsia: Lack of Correlation between  
 15 Dysmotility, Symptoms and Gastric Emptying”, Gut, (1998) 42:235-242. Tack J, et al., “Symptom Pattern and Gastric Emptying Rate Assessed by the Octanoic Acid Breath Test in Functional Dyspepsia” [abstract]. Gastroenterol, (1998) 114:A301. Cuomo R, et al., “Functional Dyspepsia Symptoms, Gastric Emptying and Satiety Provocation Test: Analysis of Relationships”, Scand J Gastroenterol, (2001)  
 20 36:1030-1036. Sarnelli G, et al., “Symptoms Associated with Impaired Gastric Emptying of Solids and Liquids in Functional Dyspepsia”, Am J Gastroenterol, (2003) 98:783-788.

## 2. Irritable Bowel Syndrome (IBS)

25 The second FGID of interest, IBS, is diagnosed when a patient’s symptoms include persistent abdominal pain or discomfort, in the absence of other explanatory organic disease, along with at least two of the following: relief of pain with defecation, onset of symptoms associated with a change in frequency of stools and/or onset of symptoms associated with a change in appearance/form of stools.  
 30 Thompson WG, et al., “Rome II: A Multinational Consensus Document on Gastrointestinal Disorders – Functional Bowel Disorders and Functional Abdominal Pain”, Gut, (1999) ;45(Suppl II):II43-II47.

In addition to colonic dysmotility, a number of other GI motility abnormalities have been identified, including delayed gastric emptying,  
 35 gastroparesis, and small intestine motility abnormalities. Vassallo MJ, et al., “Colonic Tone and Motility in Patients with Irritable Bowel Syndrome”, Mayo Clin Proc, (1992);67:725-731. Van Wijk HJ, et al., “Gastric Emptying and Dyspeptic Symptoms in the Irritable Bowel Syndrome”, Scand J Gastroenterol, (1992);27:99-

102. Evans PR, et al., "Gastroparesis and Small Bowel Dysmotility in Irritable Bowel Syndrome", Dig Dis Sci (1997);42:2087-2093. Cann PA, et al. "Irritable Bowel Syndrome: Relationship of Disorders in the Transit of a Single Solid Meal to Symptoms Patterns", Gut, (1983);24:405-411. Kellow JE, et al., "Dysmotility of the Small Intestine in Irritable Bowel Syndrome", Gut, (1988);29:1236-1243. Evans PR, et al., "Jejunal Sensorimotor Dysfunction in Irritable Bowel Syndrome: Clinical and Psychosocial Features", Gastroenterol, (1996);110:393-404. Schmidt T, et al., "Ambulatory 24-Hour Jejunal Motility in Diarrhea-Predominant Irritable Bowel Syndrome", J Gastroenterol, (1996);31:581-589. Simren M, et al., "Abnormal Propagation Pattern of Duodenal Pressure Waves in the Irritable Bowel Syndrome (IBS)", Dig Dis Sci, (2000);45:2151-2161.

A related finding is that patients with constipation-predominant IBS have evidence of decreased vagal tone, while diarrhea-predominant IBS is associated with evidence of increased sympathetic activity. Aggarwal A, et al., "Predominant Symptoms in Irritable Bowel Syndrome Correlate with Specific Autonomic Nervous system Abnormalities", Gastroenterol, (1994);106:945-950.

There is no cure for IBS. Treatments include supportive palliative care (antidiarrheals, dietary modification and counseling).

A recently approved drug to treat selected patients with FGIDs is tegaserod maleate sold under the tradename "Zelnorm<sup>®</sup>" by Novartis Pharmaceuticals Corp., East Hanover, New Jersey, USA. The product literature on Zelnorm recognizes the enteric nervous system is a key element in treating IBS. The literature suggests Zelnorm<sup>®</sup> acts to enhance basal motor activity and to normalize impaired motility. Novartis product description, Zelnorm<sup>®</sup>, July 2002 (T2002-19). Zelnorm's approved use is limited to females with constipation-related IBS. It is for short-term use only.

### B. Gastroparesis

The third disease indication discussed here, gastroparesis (or delayed gastric emptying) is associated with upper GI symptoms such as nausea, vomiting fullness, bloating and early satiety. Gastroparesis can be caused by many underlying conditions. The most important, because of chronicity and prevalence, are diabetes, idiopathic and post-surgical. Hornbuckle K, et al. "The Diagnosis and Work-Up of the Patient with Gastroparesis", J Clin Gastroenterol, (2000);30:117-124. GI dysmotility in the form of delayed gastric emptying is, by definition, present in these patients.

In patients with Type 1 diabetes mellitus and delayed gastric emptying, there appears to be a relationship between delayed gastric emptying and low vagal tone.

Merio R, et al., "Slow Gastric Emptying in Type 1 Diabetes: Relation to Autonomic and Peripheral Neuropathy, Blood Glucose, and Glycemic Control", Diabetes Care, (1997);20:419-423. A related finding is that patients with Type 1 diabetes have low vagal tone in association with increased gastric antral size, possibly contributing to the dysmotility-associated symptoms seen in these patients. Undeland KA, et al., "Wide Gastric Antrum and Low Vagal Tone in Patients with Diabetes Mellitus Type 1 Compared to Patients with Functional Dyspepsia and Healthy Individuals", Dig Dis Sci, (1996);41:9-16.

The current treatments for gastroparesis are far from satisfactory. They include supportive care, such as dietary modification, prokinetic drugs, and; when required, interventions such as intravenous fluids and placement of a nasogastric tube may be needed.

### C. Gastroesophageal Reflux Disease (GERD)

The fourth indication, GERD, can be associated with a wide spectrum of symptoms, including dyspepsia, reflux of gastric contents into the mouth, dysphagia, persistent cough, refractory hyperreactive airway disease and even chronic serous otitis media. Sontag SJ, et al., "Asthmatics with Gastroesophageal Reflux: Long Term Results of a Randomized Trial of Medical and Surgical Antireflux Therapies", Am J Gastroenterol, (2003);98:987-999. Poelmans J, et al., "Prospective Study on the Incidence of Chronic Ear Complaints Related to Gastroesophageal Reflux and on the Outcome of Antireflux Therapy", Ann Otol Rhinol Laryngol, (2002);111:933-938.

GERD is considered to be a chronic condition for which long-term medical therapy and/or surgical therapy is often deemed necessary, in significant part because esophageal adenocarcinoma is sometimes a consequence of GERD. DeVault KR, et al., "Updated Guidelines for the Diagnosis and Treatment of Gastroesophageal Reflux Disease", Am J Gastroenterol, (1999);94:1434-1442. Lagergren J, et al., "Symptomatic Gastroesophageal Reflux as a Risk Factor for Esophageal Adenocarcinoma", New Engl J Med, (1999);340:825-831.

The underlying pathophysiological mechanisms in GERD are considered to be transient lower esophageal relaxations (TLESRs) in the presence of either an inadequate pressure gradient between the stomach and the esophagus across the lower esophageal sphincter and/or low amplitude esophageal activity at times when gastric contents do reflux into the esophagus. In addition, gastric distention is thought to be associated with an increase in TLESRs. Mittal RK, et al., "Mechanism of Disease: The Esophagogastric Junction", New Engl J Med, (1997);336:924-932. Scheffer RC, et al., "Elicitation of Transient Lower Oesophageal Sphincter

Relaxations in Response to Gastric Distension”, Neurogastroenterol Motil. (2002);14:647-655.

GERD is generally considered to be the result of a motility disorder which permits the abnormal and prolonged exposure of the esophageal lumen to acidic gastric contents. Hunt, “The Relationship Between The Control Of pH And Healing And Symptom Relief In Gastro-Oesophageal Reflux Disease”, Ailment Pharmacol Ther., 9 (Suppl. 1) pp. 3 – 7 (1995). Many factors are believed to contribute to the onset of GERD. These include transient lower esophageal sphincter relaxations (as previously described), decreased LES resting tone, delayed stomach emptying and an ineffective esophageal clearance.

Certain drugs have had some effectiveness at controlling GERD but fail to treat underlying causes of the disease. Examples of such drugs are H<sub>2</sub>-receptor antagonists (which control gastric acid secretion in the basal state) and proton pump inhibitors (which control meal-stimulated acid secretion). Hunt, id. Both classes of drugs can raise intragastric pH to or about 4 for varying durations. Hunt, supra.

Surgery treatments are also employed for the treatment of GERD and include techniques for bulking the lower esophageal sphincter such as fundoplication and techniques described in US Pat. No. 6,098,629 Johnson et al, Aug 8, 2000. Other surgical techniques include placement of pacemakers for stimulating muscle contractions in the esophageal sphincter, the stomach muscles or in the pyloric valve. U.S. Pat. No. 6,104,955 to Bourgeois, U.S. Pat. No. 5,861,014 to Familoni.

A summary of GERD treatments can be found in DeVault, et al., “Updated Guidelines for the Diagnosis and Treatment of Gastroesophageal Reflux Disease”, Amer. J. of Gastroenterology, Vol. 94, No. 6, pp. 1434 – 1442 (1999).

Notwithstanding multiple attempts at various types of treatment, GERD continues to be a serious disease proving to be difficult to treat by any of the foregoing prior art techniques. In view of the foregoing and notwithstanding various efforts exemplified in the prior art, there remains a need for an effective treatment for GERD. It is an object of the present invention to provide a novel treatment and novel apparatus for the treatment of GERD.

#### **D. Electrical Stimulation to Treat GI Disorders**

Treatment of gastrointestinal diseases through nerve stimulation have been suggested. For example, U.S. Pat. No. 6,238,423 to Bardy dated May 29, 2001 describes a constipation treatment involving electrical stimulation of the muscles or related nerves of the gut. U.S. Pat. No. 6,571,127 to Ben-Haim et al. dated May 27, 2003 describes increasing motility by applying an electrical field to the GI tract. U.S. Pat. No. 5,540,730 to Terry, Jr. et al., dated July 30, 1996 describes a motility

treatment involving vagal stimulation to alter GI contractions in response to a sense condition indicative of need for treatment. The '730 patent also uses a definition of dysmotility more restrictive than in the present application. In the '730 patent, dysmotility is described as hyper- or hypo-contraction. In the present application, dysmotility is a broader concept to refer to all abnormalities of gastric emptying or bowel transfer regardless of cause. U.S. Pat. No. 6,610,713 to Tracey dated August 26, 2003 describes inhibiting release of a proinflammatory cytokine by treating a cell with a cholinergic agonist by stimulating efferent vagus nerve activity to inhibit the inflammatory cytokine cascade.

A substantial body of literature is developed on nerve stimulation. For example, in Dapoigny et al., "Vagal influence on colonic motor activity in conscious nonhuman primates", Am. J. Physiol., 262: G231 – G236 (1992), vagal influence on colonic motor activity was investigated in conscious monkeys. To block antidromic interference, the vagus was blocked via vagal cooling and a vagal stimulation electrode was implanted distal to the vagal block. It was noted that vagal efferent stimulation increased contractile frequency and that the vagus has either a direct or indirect influence on fasting and fed colonic motor activity throughout the colon, and that a non-adrenergic, noncholinergic inhibitory pathway is under vagal control.

Colonic and gastric stimulation are also described in a number of articles associated with M. P. Mintchev. These include: Mintchev, et al., "Electrogastrographic impact of multi-site functional gastric electrical stimulation", J. of Medical Eng. & Tech., Vol. 23, No. 1 pp. 5 – 9 (1999); Rashev, et al., "Three-dimensional static parametric modeling of phasic colonic contractions for the purpose of microprocessor-controlled functional stimulation", J. of Medical Eng. & Tech., Vol. 25, No. 3 pp. 85 – 96 (2001); Lin et al., "Hardware – software co-design of portable functional gastrointestinal stimulator system", J. of Medical Eng. & Tech., Vol. 27, No. 4 pp. 164 – 177 (2003); Amaris et al., "Microprocessor controlled movement of solid colonic content using sequential neural electrical stimulation", Gut, 50: pp 475 – 479 (2002) and Rashev et al., "Microprocessor-Controlled Colonic Peristalsis", Digestive Diseases and Sciences, Vol. 47, No. 5, pp. 1034 – 1048 (2002).

The foregoing references describe nerve stimulation to stimulate muscular contraction in the GI tract. As will be more fully discussed, the present invention utilizes vagal stimulation to improve vagal tone (similar in concept to improving cardiac electrical tone through cardiac pacing) and/or to treat GI disorders by altering the nature of duodenum contents by stimulation pancreatic and biliary



output. The invention is also applicable to treating other diseases such as neuropsychiatric disorders.

Vagal tone has been shown to be associated with dyspepsia. Hjelland, et al., "Vagal tone and meal-induced abdominal symptoms in healthy subjects", Digestion, 65: 172 – 176 (2002). Also, Hausken, et al., "Low Vagal Tone and Antral Dysmotility in Patients with Functional Dyspepsia", Psychosomatic Medicine, 55: 12 – 22 (1993). Also, decreased vagal tone has been associated with irritable bowel syndrome. Heitkemper, et al., "Evidence for Automatic Nervous System Imbalance in Women with Irritable Bowel Syndrome", Digestive Diseases and Sciences, Vol. 43, No. 9, pp. 2093 – 2098 (1998).

Also, as will be discussed, the present invention includes, in several embodiments, a blocking of a nerve (such as the vagal nerve) to avoid antidromic influences during stimulation. Cryogenic nerve blocking of the vagus is described in Dapoigny et al., "Vagal influence on colonic motor activity in conscious nonhuman primates", Am. J. Physiol., 262: G231 – G236 (1992). Electrically induced nerve blocking is described in Van Den Honert, et al., "Generation of Unidirectionally Propagated Action Potentials in a Peripheral Nerve by Brief Stimuli", Science, Vol. 206, pp. 1311 – 1312. An electrical nerve block is described in Solomonow, et al., "Control of Muscle Contractile Force through Indirect High-Frequency Stimulation", Am. J. of Physical Medicine, Vol. 62, No. 2, pp. 71 – 82 (1983) and Petrofsky, et al., "Impact of Recruitment Order on Electrode Design for Neural Prosthetics of Skeletal Muscle", Am. J. of Physical Medicine, Vol. 60, No.5, pp. 243 – 253 (1981). A neural prosthesis with an electrical nerve block is also described in U.S. Patent Application Publication No. US 2002/0055779 A1 to Andrews published May 9, 2002. A cryogenic vagal block and resulting effect on gastric emptying are described in Paterson CA, et al., "Determinants of Occurrence and Volume of Transpyloric Flow During Gastric Emptying of Liquids in Dogs: Importance of Vagal Input", Dig Dis Sci, (2000);45:1509-1516.

### SUMMARY OF THE INVENTION

A method and apparatus are disclosed for treating at least one of a plurality of disorders of a patient associated with vagal activity innervating at least one of a plurality of organs of the patient at an innervation site. The method includes positioning a neurostimulator carrier within a body lumen of the patient. An electrode disposed on the carrier is positioned near a mucosal layer of the lumen. An electrical signal is applied to the electrode to modulate vagal activity by an amount selected to treat the disorder. The signal may be a blocking or a stimulation signal.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

Fig. 1 is a schematic representation of a gastric-emptying feedback loop with a patient-controlled stimulator for stimulating an organ of the loop;

5        Fig. 2 is a view similar to Fig. 1 with an automatic controller replacing the patient-controller of Fig. 1 and with feedback circuits to the automatic controller schematically represented;

10       Fig. 3 is a schematic illustration of an alimentary tract (GI tract plus non-GI organs such as the pancreas and liver) and its relation to vagal and enteric innervation;

      Fig. 4 is the view of Fig. 3 showing the application of a pacing electrode according to an embodiment of the present invention;

      Fig. 5 is a schematic representation of pacing system;

15       Fig. 6 is the view of Fig. 4 showing the application of a nerve conduction block electrode proximal to the pacing electrode;

      Fig. 7 is the view of Fig. 6 showing the application of a nerve conduction block electrode distal to the pacing electrode;

      Fig. 8 is the view of Fig. 3 showing the application of a nerve conduction block electrode according to an embodiment of the present invention;

20       Fig. 9 is a schematic representation of a patients' stomach shown partially in section and illustrating a representative placement of anterior and posterior vagus nerves with respect to the anatomy of the stomach and diaphragm;

      Fig. 10 is the view of Fig. 9 showing a further embodiment of the present invention in utilizing electrode bands;

25       Fig. 11 is a perspective view of a band for use in the embodiment of Fig. 10;

      Fig. 12 is a side sectional view of a patients' stomach in illustrating a yet alternative embodiment of the present invention;

      Fig. 13 is a side elevation view of a balloon portion of an apparatus for use in the embodiment of Fig. 12;

30       Fig. 14 is a side elevation view of an alternative embodiment of a balloon portion of an apparatus for use in the embodiment of Fig. 12;

      Fig. 15 is a side sectional view of a patients' stomach in illustrating a yet alternative embodiment of the invention of Fig. 12;

35       Fig. 16 is a side sectional view of a patients' stomach in illustrating a still further alternative embodiment of the invention of Fig. 12;

      Fig. 17 is a schematic view of a balloon with magnetic coils;

      Fig. 18 is a side sectional view of the esophagus and stomach junction and illustrating a yet further embodiment of the present invention; and

Fig. 19 is a view similar to that of Fig. 15 and illustrating an alternative embodiment of the invention shown in Fig. 15.

### **DESCRIPTION OF THE PREFERRED EMBODIMENT**

5 With reference now to the various drawing figures in which identical elements are numbered identically throughout, a description of the preferred embodiment of the present invention will now be described.

#### **A. Invention of Parent Application**

10 Figs 1 and 2 and the description which follow are from the aforementioned U.S. patent application Ser. No. 10/358,093 filed February 3, 2003 filed February 3, 2003 and entitled "Method and Apparatus for Treatment of Gastroesophageal Disease (GERD)".

15 With initial reference to Fig. 1, a gastric emptying feedback loop is shown schematically for ease of illustration. The feedback loop illustrates a patient's stomach S which is provided with food from the esophagus E. A lower esophageal sphincter LES is shown positioned between the esophagus E and the stomach S. The lower esophageal sphincter normally provides control of reflux of stomach contents into the esophagus E.

20 On a proximal or lower end of the stomach S the stomach discharges into the superior duodenum D which is an upper portion of the intestines. The superior duodenum D and the stomach S are separated by a pyloric valve PV which opens to permit gastric emptying from the stomach into the duodenum D.

25 Also schematically illustrated in Fig. 1 are nerve paths N providing signal flow paths from both the superior duodenum D and the stomach S to the brain B. An efferent Vagal nerve VN connects the brain B to the pancreas P of the patient. A conduit (pancreatic duct PD) extends from the pancreas P and discharges into the superior duodenum D.

30 The presence of food contents within the duodenum D (such contents being referred to as "chyme") may prevent passage of gastric content of the stomach S past the pyloric valve PV into the duodenum D. As long as such gastric contents cannot be passed into the duodenum D, such contents can be forced retrograde past the lower esophageal sphincter LES and into the esophagus E creating the symptoms and discomfort of GERD. The contents discharging from the stomach S into the  
35 duodenum D are acidic (and high osmolality) and reside in the duodenum D until pH is elevated (close to a neutral pH of 6 – 7) and osmolality is normalized.

The elevation of pH and reduction of osmolality of chyme in the duodenum D results from exocrine secretion being administered from the pancreas P and from

bile from the liver into the duodenum D. This raises the pH and lowers the osmolality of the duodenum D content permitting discharge from the duodenum D and thereby permitting gastric emptying across the pyloric valve PV.

According to the present invention gastroesophageal reflux disease (GERD) results from a derangement of the feedback loops involved in upper GI digestion and motility control. This problem encompasses receptors and reflexes that regulate the propulsive contractions of the stomach, upper duodenum and biliary tree and the secretions of the exocrine pancreas. The interaction of these receptors and reflexes control gastric emptying (by coordinating gastric propulsive contractions and sphincter [primarily pyloric] tone) and regulate the pH and osmolality of the chyme in the duodenum. This chemo-regulation is mediated through control of bile delivery and stimulation of secretion by the exocrine pancreas of fluid delivered to the superior duodenum. Chey et al., "Neural Hormonal Regulation of Exocrine Pancreatic Secretion", *Pancreatology*, pp. 320 – 335 (2001).

Normally, ingestate delivered to the stomach is mixed by low intensity gastric mixing contractions with the enzymatic, ionic, including hydrogen ion ( $H^+$ ), and water secretions of the glands of the stomach. When the material is adequately reduced in size and is a smooth consistency, the fluid, now called chyme, is delivered to the ampulla of the small intestine by the much stronger propulsive, or emptying, contractions of the stomach coupled with transitory relaxation of the pyloric sphincter. This material is at a very low pH (about 2) and high osmolality, which activates receptors, including those for  $H^+$  and osmotic pressure, which are abundant in the wall of the ampulla. This receptor activation initiates the series of reflexes that cause pancreatic exocrine secretion to be delivered into the superior duodenum and ampulla. This fluid contains digestive enzymes, water and buffering compounds to raise the pH, and reduce the osmolality, of the chyme.

Once a neutral pH and physiological osmolality are achieved, then propulsive contractions in the superior duodenum move the chyme out of the superior portion into the length of the duodenum; At which point the stretch and baro-receptors in the ampulla allow the pyloric sphincter to relax and another bolus of gastric contents is delivered into the ampulla by the peristaltic gastric emptying contractions. This material, at a very low pH (less than 2), activates hydrogen ion ( $H^+$ ) on receptors of the ampulla (upper most portion of the duodenum) causing the pancreatic fluids to be delivered to the material in the ampulla restarting the cycle as described above. Chapter 3, "The Stomach", *Gastrointestinal System*, 2<sup>nd</sup> Ed., M.S. Long editor, Mosby Publisher, London (2002).

If the control system is down regulated by, for example, by increased pH of gastric contents entering the ampulla, feedback may thereby be reduced from the  $H^+$

receptors in the duodenum that stimulate pancreatic exocrine secretion and bile delivery to the duodenum, then movement of chyme from the superior duodenum is delayed, causing delay of gastric emptying. Mabayo, et al., "Inhibition of Food Passage by Omeprazole in the Chicken", European J. of Pharmacology, pp. 161 – 165 (1995).

In GERD, this reflex is inhibited in such a way that the stomach empties more slowly so that the gastric emptying contractions force gastric contents to flow retrograde into the esophagus. This is a result of the situation in which the gastric emptying contractions are vigorous but must operate against a contracted pyloric sphincter. These vigorous peristaltic contractions eventually begin to force gastric contents to flow retrograde into the esophagus because of the inherent imbalance between a very strong pyloric sphincter and a much weaker gastroesophageal sphincter. The delay in gastric emptying is directly related to a slow down in the transport of chyme out of the ampulla and superior duodenum. The drugs used to treat this disease raise pH further dampening the hydrogen-receptor-pancreatic secretion loop, further delaying gastric emptying. Benini, "Gastric Emptying and Dyspeptic Symptoms in Patients with Gastroesophageal Reflux", Amer. J. of Gastroenterology, pp. 1351 – 1354 (1996).

The present invention is directed towards reestablishing the link between gastric emptying and pancreatic secretion delivery, thereby addressing the main pathology of this disease by shortening chyme residence time in the superior duodenum so that intestinal contents move into the distal digestive tract in a more normal manner. According to a first embodiment, this is done by stimulating the H<sup>+</sup> ion receptors or by stimulation of the pancreas directly or via its para-sympathetic innervation (pre-ganglionic Vagal nerves). Stimulation of pancreatic exocrine secretion has been shown by direct stimulation of the thoracic vagus nerves in dogs. Kaminski et al., "The Effect of Electrical Vagal Stimulation on Canine Pancreatic Exocrine Function", Surgery, pp. 545 – 552 (1975). This results in a more rapid (normal) neutralization of chyme in the ampulla, allowing it move down the duodenum more quickly so that gastric emptying is returned to a more normal pace.

Acidity (pH) can be assessed by measuring bicarbonate. It will be understood that references to -H includes such indirect measurements. Also, effects of the therapy described herein can be assessed and/or controlled by measuring an indication of pancreatic exocrine secretion or bile (e.g., HCO<sub>3</sub><sup>-</sup>).

An alternative embodiment uses gastrocopic delivery of a paralyzing agent (e.g. botulism toxin) to the pyloric valve along with use of H<sub>2</sub> antagonists or PPI's to manage the acidity of the chyme reaching the duodenum.

As an additional alternative to pancreatic stimulation, the gall bladder can be stimulated to encourage bile movement into the duodenum. Shown schematically in the figures, the gall bladder GB resides below the liver L. The gall bladder is connected to the small intestine (specifically the duodenum D) via a bile duct BD.

5 The bile duct BD can discharge directly into the duodenum D or via the pancreatic duct PD as shown. The bile can normalize the chyme to accelerate duodenal emptying. Bile consists of bile acids (detergents that emulsify lipids), cholesterol, phospholipids, electrolytes such as ( $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{+2}$ ,  $\text{Cl}^-$ ,  $\text{HCO}_3^-$ ) and  $\text{H}_2\text{O}$ . Chapter 4, "The Liver and Biliary Tract", Gastrointestinal System, 2<sup>nd</sup> Ed., M.S. Long editor,

10 Mosby Publisher, London (2002). The gall bladder GB or bile duct can be stimulated indirectly via stimulation of the vagal nerve VN or directly stimulated by an electrode 11 (shown in phantom lines).

As illustrated in the figures, an electrical stimulator 10, 20 which may be implanted is provided which alternatively may be directly connected to the Vagal

15 nerve VN or the pancreas P to stimulate the pancreas directly or indirectly to excrete exocrine into the duodenum D (or more distally into the small intestine – e.g., into the jejunum) and increase the pH of chyme in the duodenum D as described. Alternatively, the same can be done to promote bile release. The frequency may be varied to maximize the response and selectively stimulate exocrine instead of

20 endocrine secretions. Rösch et al., "Frequency-Dependent Secretion of Pancreatic Amylase, Lipase, Trypsin, and Chymotrypsin During Vagal Stimulation in Rats", Pancreas, pp. 499 – 506 (1990). See, also, Berthoud et al., "Characteristics of Gastric and Pancreatic Responses to Vagal Stimulation with Varied Frequencies: Evidence for Different Fiber Calibers?", J. Auto. Nervous Sys., pp. 77 – 84 (1987)

25 (showed frequency-response relationship with insulin, i.e., significantly less insulin was released at lower frequencies – 2 Hz v. 8 Hz – also, frequency-response curves evidenced distinctly different profiles for gastric, pancreatic and cardiovascular responses.) Slight insulin release can maximize pancreatic exocrine secretion. Chey et al., "Neural Hormonal Regulation of Exocrine Pancreatic Secretion",

30 Pancreatology, pp. 320 – 335 (2001).

With a patient control stimulation as shown in Fig. 1, the patient may activate the stimulator 10 by remote transmitter to stimulate an electrical charge either after eating (e.g., about 60 to 90 minutes after eating) or on onset of GERD symptoms. It will be appreciated that there are a wide variety of nerve stimulators

35 and organ stimulators available for implantation and are commercially available and which include connectors for connecting directly to nerves.

Fig. 2 illustrates an additional embodiment where the patient activated loop is replaced with an automatic loop having a programmable stimulator 20 which

receives as an input signals from sensors in the duodenum to measure pH, osmolality or strain (e.g., from baro-sensors) on the duodenum indicating filling or may measure acidity in the esophagus or strain on the lower esophageal sphincter LES or stomach S all of which may be provided to the implantable controller 20 which can be provided with desirable software to process the incoming signals and generate a stimulating signal to either the vagal nerve, the pancreas P or the duodenum D (or jejunum) directly in response to such received signals. It will be appreciated that stimulators and controllers are well within the skill of the art. U.S. Pat. No. 5,540,730 teaches a neurostimulator to stimulate a vagus nerve to treat a motility disorder. U.S. Pat. No. 5,292,344 teaches gastrointestinal sensors, including pH sensors.

### **B. Application of Parent Application to Treatments Other than GERD**

In addition to treatment of GERD, the foregoing invention is applicable to treatment of a plurality of GI diseases associated with delayed gastric emptying or altered autonomic activity. These include functional gastrointestinal disorders and gastroparesis. Furthermore, applicants have determined that duodenal content impacts a plurality of motility disorders throughout the bowels and can diseases associated with dysmotility (e.g., irritable bowel syndrome). Accordingly it is an object of the present invention to use the teachings of the aforementioned parent application to treat GI disorders associated with delayed gastric emptying and abnormal intestinal transport.

### **C. Additional Disclosure of the Present Application**

#### **1. Enteric Innervation**

Fig. 3 is a schematic illustration of an alimentary tract (GI tract plus non-GI organs such as the pancreas and gall bladder, collectively labeled PG) and its relation to vagal and enteric innervation. The lower esophageal sphincter (LES) acts as a gate to pass food into the stomach S and, assuming adequate function of all components, prevent reflux. The pylorus PV controls passage of chyme from the stomach S into the intestines I (collectively shown in the figures and including the large intestine or colon and the small intestine including the duodenum, jejunum and ileum).

The biochemistry of the contents of the intestines I is influenced by the pancreas P and gall bladder PG which discharge into the duodenum. This discharge is illustrated by dotted arrow A.

The vagus nerve VN transmits signals to the stomach S, pylorus PV, pancreas and gall bladder PG directly. Originating in the brain, there is a common

vagus nerve VN in the region of the diaphragm (not shown). In the region of the diaphragm, the vagus VN separates into anterior and posterior components with both acting to innervate the GI tract. In Figs. 3, 5 – 8, the anterior and posterior vagus nerves are not shown separately. Instead, the vagus nerve VN is shown

5 schematically to include both anterior and posterior nerves.

The vagus nerve VN contains both afferent and efferent components sending signals away from and to, respectively, its innervated organs.

In addition to influence from the vagus nerve VN, the GI and alimentary tracts are greatly influenced by the enteric nervous system ENS. The enteric

10 nervous system ENS is an interconnected network of nerves, receptors and actuators throughout the GI tract. There are many millions of nerve endings of the enteric nervous system ENS in the tissues of the GI organs. For ease of illustration, the enteric nervous system ENS is illustrated as a line enveloping the organs innervated by the enteric nervous system ENS

15 The vagus nerve VN innervates, at least in part, the enteric nervous system ENS (schematically illustrated by vagal trunk VN3 which represents many vagus-ENS innervation throughout the cut). Also, receptors in the intestines I connect to the enteric nervous system ENS. Arrow B in the figures illustrates the influence of duodenal contents on the enteric nervous system ENS as a feedback to the secretion

20 function of the pancreas, liver and gall bladder. Specifically, receptors in the intestine I respond the biochemistry of the intestine contents (which are chemically modulated by the pancreao-biliary output of Arrow A). This biochemistry includes pH and osmolality.

In the figures, vagal trunks VN1, VN2, VN4 and VN6 illustrate

25 schematically the direct vagal innervation of the GI organs of the LES, stomach S, pylorus PV and intestines I. Trunk VN3 illustrates direct communication between the vagus VN and the ENS. Trunk VN5 illustrates direct vagal innervation of the pancreas and gall bladder. Enteric nerves ENS1 – ENS4 represent the multitude of enteric nerves in the stomach S, pylorus PV, pancreas and gall bladder PG and

30 intestines I.

While communicating with the vagus nerve VN, the enteric nervous system ENS can act independently of the vagus and the central nervous system. For example, in patients with a severed vagus nerve (vagusotomy – an historical procedure for treating ulcers), the enteric nervous system can operate the gut. Most enteric

35 nerve cells are not directly innervated by the vagus. Gershon, “The Second Brain”, Harper Collins Publishers, Inc, New York, NY p. 19 (1998)

In Fig. 3, the vagus VN and its trunks (illustrated as VN1 – VN6) and the enteric nervous system ENS are shown in phantom lines to illustrate reduced vagal



and enteric nerve tone (i.e., sub-optimal nerve transmission levels). Reduced vagal and enteric tone contribute directly to the ineffectiveness of the GI organs as well as indirectly (through reduced pancreatic/biliary output). The reduced pancreatic/biliary output is illustrated by the dotted presentation of arrow A. As previously discussed, the vagus can be stimulated to stimulate pancreatic or biliary output. Therefore, the reduced output of arrow A results in a reduced feedback illustrated by the dotted presentation of arrow B.

## 2. Enteric Rhythm Management (ERM)

The benefits of the present invention are illustrated in Fig. 4 where a stimulating or pacing electrode PE is applied to the vagus VN. While only one electrode is shown in Fig. 4, separate electrodes could be applied to both the anterior and posterior vagus nerves (or to the common vagus or vagal branches). In a preferred embodiment, the electrode PE is placed a few centimeters below the diaphragm and proximal to stomach and pancreo/biliary innervation. While this placement is presently preferred for ease of surgical access, other placement locations may be used.

By pacing the vagus through the pacing electrode, vagal tone is optimized by either up- or down-regulation. With reference to the parasympathetic and enteric nervous systems, "tone" refers to basal activity of a nerve or nervous system facilitating appropriate physiologic response to a patient's internal environment. For example, low vagal tone implies a reduction in vagus nerve activity resulting in decreased response of the alimentary tract to ingested food. As used in the present application, "pacing" is not limited to mean timed events coordinated with specifically timed physiologic events. Instead, pacing means any electrical stimulation of a nerve trunk to induce bi-directional propagation of nervous impulses in the stimulated nerve.

The operating effectiveness of the vagus is enhanced so that local physiological signals generated in the enteric nervous system (or sent to the brain from the organs) are more appropriately responded to within the alimentary tract. Due to its innervation of the enteric nervous system, pacing of the vagus enhances the functional tone of the enteric nervous system. By enhancing the functional tone it will be noted that the stimulation pacing is elevating the degree of functionality of the vagus and enteric nerves. In this context, "pacing" is not meant to mean timed pulsed coordinated with muscular contractions or synchronized with other events. Pacing means elevating the activity level of the nerves.

Tonal enhancement of the vagus and enteric nerves is illustrated by the solid lines for the nerves VN, ENS in Fig. 4. Vagal trunk VN5 is in solid line to illustrate

enhanced tone of the many vagal nerve components communicating with the enteric nervous system ENS. Direct vagal innervation of the LES, stomach S, pylorus PV and intestines I remains shown as low tone indicated by phantom lines VN1, VN2, VN4, VN6. The tonal pacing described herein is not intended to trigger or drive the muscular contractility of these organs. The stimulation is not intended to be timed to trigger contractility and is not provided with an energy level sufficient to drive peristaltic contractions. Instead, these functions remain controlled by the central and enteric nerves systems. The enhanced nerve tone provided by the present invention permits these functions to occur.

Pacing to enhance vagal tone is not initiated in response to any senses event (or in anticipation of an immediate need to GI activity). Instead, the pacing can be done intermittently over the day to provide an enhanced level of operating functionality to the vagus. By way of non-limiting example, the stimulation pacing can be done during awake hours. For example, every ten minutes, pacing signals can be sent to the pacing electrodes. The pacing signals have a duration of 30 seconds with a current of 4mA, a frequency of 12 Hz and an impulse duration of 2 msec. These parameters are representative only. A wide range of signal parameters may be used to stimulate the vagus nerve. Examples of these are recited in the afore-referenced literature

As will be further discussed, the present invention permits ERM to be uniquely designed and modified by an attending physician to meet the specific needs of individual patients. For example, pacing can be limited to discrete intervals in the morning, afternoon and evening with the patient free to coordinate meals around these events.

In addition to enhancing vagal and enteric tone directly, the pacing also enhances the pancreatic and biliary output for the reasons discussed above. Namely, while ERM does not drive muscular events over nerve trunks VN1, VN2, VN4, VN6, the enhanced tone stimulates pancreo-biliary output over trunk VN5 (illustrated by the solid line of VN5 in Fig. 4). This enhanced output is illustrated as solid arrow A' in Fig. 4. As a consequence there is a greater feedback to the intestinal receptors as illustrated by solid arrow B' in Fig. 4. This enhanced biochemistry feedback further enhances the tone of the enteric nervous system ENS.

### 3. Implantable Pacing Circuit

A representative pacing circuit 100 is schematically shown in Fig. 5. Similar to cardiac pacing devices, an implantable controller 102 contains an induction coil 104 for inductive electrical coupling to a coil 106 of an external controller 108. The implantable controller 102 includes anterior and posterior pulse generators 110, 112

electrically connected through conductors 114, 116 to anterior and posterior pacing electrodes 118, 120 for attachment to anterior and posterior trunks, respectively, of the vagus nerve VN. The implantable controller 102 also includes a battery 122 and a CPU 124 which includes program storage and memory. The timing and  
5 parameters of the pulse at the electrodes 118, 120 can be adjusted by inductively coupling the external controller 108 to the implantable controller 102 and inputting pacing parameters (e.g., pulse width, frequency and amplitude).

While a fully implantable controller 102 is desirable, it is not necessary. For example, the electrodes 118, 120 can be implanted connected to a receiving antenna  
10 placed near the body surface. The control circuits (i.e., the elements 124, 110, 112 and 108) can be housed in an external pack worn by the patient with a transmitting antenna held in place on the skin over the area of the implanted receiving antenna. Such a design is forward-compatible in that the implanted electrodes can be later substituted with the implantable controller 102 at a later surgery if desired.

15 Although not shown in Fig. 5, the controller 102 can also include circuits generating nerve conduction block signals (as will be described) which connect to electrodes which may be positioned on a nerve proximally, distally (or both) of the electrodes 118, 120.

#### 20 4. Nerve Conduction Block

Fig. 6 shows an alternative embodiment using a nerve conduction blocking electrode PBE proximal to the pacing electrode for providing a conduction block. A nerve block is, functionally speaking, a reversible vagotomy. Namely, application of the block at least partially prevents nerve transmission across the site of the block.  
25 Removal of the block restores normal nerve activity at the site. A block is any localized imposition of conditions that at least partially diminish transmission of impulses.

The vagal block may be desirable in some patients since unblocked pacing may result in afferent vagal and antidromic efferent signals having undesired effect  
30 on organs innervated by the vagus proximal to the GI tract (e.g., undesirable cardiac response). Further, the afferent signals of the pacing electrode PE can result in a central nervous system response that tends to offset the benefits of the pacing electrode on the ENS and pancreo/biliary function, thereby reducing the GI and enteric rhythm management effectiveness of vagal pacing.

35 The block may be intermittent and applied only when the vagus is paced by the pacing electrode PE. The preferred nerve conduction block is an electronic block created by a signal at the vagus by an electrode PBE controlled by the implantable controller (such as controller 102 or an external controller). The nerve

conduction block can be any reversible block. For example, cryogenics (either chemically or electronically induced) or drug blocks can be used. An electronic cryogenic block may be a Peltier solid-state device which cools in response to a current and may be electrically controlled to regulate cooling. Drug blocks may include a pump-controlled subcutaneous drug delivery.

With such an electrode conduction block, the block parameters (signal type and timing) can be altered by a controller and can be coordinated with the pacing signals to block only during pacing. A representative blocking signal is a 500Hz signal with other parameters (e.g., timing and current) matched to be the same as the pacing signal. While an alternating current blocking signal is described, a direct current (e.g., -70mV DC) could be used. The foregoing specific examples of blocking signals are representative only. Other examples and ranges of blocking signals are described in the afore-mentioned literature. For example, the nerve conduction block is preferably within the parameters disclosed in Solomonow, et al., "Control of Muscle Contractile Force through Indirect High-Frequency Stimulation", Am. J. of Physical Medicine, Vol. 62, No. 2, pp. 71 – 82 (1983). Particularly, the nerve conduction block is applied with electrical signal selected to block the entire cross-section of the nerve (e.g., both afferent, efferent, myelinated and nonmyelinated fibers) at the site of applying the blocking signal (as opposed to selected sub-groups of nerve fibers or just efferent and not afferent or visa versa) and, more preferably, has a frequency selected to exceed the 200 Hz threshold frequency described in Solomonow et al. Further, preferred parameters are a frequency of 500 Hz (with other parameters, as non-limiting examples, being amplitude of 4 mA, pulse width of 0.5 msec, and duty cycle of 5 minutes on and 10 minutes off). As will be more fully described, the present invention gives a physician great latitude in selected pacing and blocking parameters for individual patients.

Similar to Fig. 4, the vagus VN and enteric nervous system ENS in Fig. 6 distal to the block PBE are shown in solid lines to illustrate enhanced tone (except for the direct innervation VN1, VN2, VN4, VN6 to the GI tract organs). Similarly, arrows A', B' are shown in solid lines to illustrate the enhanced pancreo-biliary output and resultant enhanced feedback stimulation to the enteric nervous system ENS. The proximal vagus nerve segment VNP proximal to the block PBE is shown in phantom lines to illustrate it is not stimulated by the pacing electrode PE while the blocking electrode PBE is activated.

## 5. Proximal and Distal Blocking

Fig. 7 illustrates the addition over Fig. 6 of a nerve conductive block DBE distal to the pacing electrode PE. The proximal block PBE prevents adverse events resulting from afferent signals and heightens the GI effectiveness by blocking antidromic interference as discussed with reference to Fig. 6.

In Fig. 7, the distal block DBE is provided in the event there is a desire to isolate the pacing effect of electrode PE. For example, a physician may wish to enhance the vagus and enteric activity in the region proximal to the duodenum but may wish to avoid stimulating pancreo-biliary output. For example, a patient may have a GI problem without apparent colon dysfunction (e.g., gastroparesis functional dyspepsia without bowel symptoms). Placing the distal block DBE on a branch of the vagus between the pacing electrode PE and the pancreas and gall bladder PG prevents increased pancreo-biliary output and resultant feedback (illustrated by dotted arrows A and B in Fig. 7 and dotted distal vagal nerve segment VND and vagal trunk VN5).

## 6. Blocking As An Independent Therapy

Fig. 8 illustrates an alternative embodiment of the invention.

In certain patients, the vagus nerve may be hyperactive contributing to diarrhea-dominant IBS. Use of a blocking electrode alone in the vagus permits down-regulating the vagus nerve VN, the enteric nervous system ENS and pancreo-biliary output. The block down-regulates efferent signal transmission. In Fig. 8, the hyperactive vagus is illustrated by the solid line of the proximal vagus nerve segment VNP. The remainder of the vagus and enteric nervous system are shown in reduced thickness to illustrate down-regulation of tone. The pancreo-biliary output (and resulting feedback) is also reduced. In Fig. 8, the blocking electrode BE is shown high on the vagus relative to the GI tract innervation (e.g., just below the diaphragm), the sole blocking electrode could be placed lower (e.g., just proximal to pancreo/biliary innervation VN5). Blocking of the entire vagus as described above can be used to down-regulate the vagus for various benefits including: pancreatitis and obesity treatments. Further, blocking the vagus interrupts the vagally-mediated neurogenic inflammatory arc.

The use of blocking as an independent therapy also permits treatment for pancreatitis by down regulating vagal activity and pancreatic output including pancreatic exocrine secretion. Also, the blocking may be used as a separate treatment for reducing discomfort and pain associated with gastrointestinal disorders or other vagally mediated pain (i.e., somatic pain sensations transmitted along any nerve fibers with pain sensation modulated by vagal afferent fibers). A nerve

stimulation to treat pain is described in U.S. patent application publication No. US2003/0144709 to Zabara et al., published July 31, 2003.

It will be appreciated that patient discomfort and pain is a primary complaint associated with many gastrointestinal disorders. As used in the present application (and appended claims), it will be appreciated that a treatment of a gastrointestinal disorder may include a treatment of a patient's perception of pain without any additional functional therapy associated with a gastrointestinal disorder. Vagal blocking as described herein can treat gastrointestinal pain or discomfort (including that associated with Crohn's disease) and chronic somatic pain as well as the inflammatory basis of Crohn's disease. The vagal blocking as described herein can also treat nausea secondary, for example, to chronic cancer chemotherapy.

### 7. Application to Obesity

The foregoing discussion has been described in a preferred embodiment of treating FGIDs, gastroparesis and GERD. Obesity is also treatable with the present invention.

Recent literature describes potential obesity treatments relative to gut hormone fragment peptide YY<sub>3-36</sub>. See, e.g., Batterham, et al., "Inhibition of Food Intake in Obese Subjects by Peptide YY<sub>3-36</sub>", New England J. Med., pp. 941 – 948 (September 4, 2003) and Korner et al., "To Eat or Not to Eat – How the Gut Talks to the Brain", New England J. Med., pp. 926 – 928 (September 4, 2003). The peptide YY<sub>3-36</sub> (PPY) has the effect of inhibiting gut motility through the phenomena of the ileal brake. Vagal afferents create a sensation of satiety.

The present invention can electrically simulate the effects of PPY by using the vagal block to down-regulate afferent vagal activity to create a desired sensation of satiety. Since the down-regulation does not require continuous blocking signals, the beneficial efferent signals are permitted.

### 8. Application to Other Therapies

There are numerous suggestions for vagal pacing or stimulation to treat a wide variety of diseases. For example, U.S. Pat. No. 5,188,104 dated February 23, 1993 describes vagal stimulation to treat eating disorders. U.S. Pat. No. 5,231,988 dated August 3, 1993 describes vagal stimulation to treat endocrine disorders. U.S. Pat. No. 5,215,086 dated June 1, 1993 describes vagal stimulation to treat migraines. U.S. Pat. No. 5,269,303 dated December 14, 1993 describes vagal stimulation to treat dementia. U.S. Pat. No. 5,330,515 dated July 19, 1994 describes vagal stimulation to treat pain. U.S. Pat. No. 5,299,569 dated April 5, 1994 describes vagal stimulation to treat neuropsychiatric disorders. U.S. Pat. No. 5,335,657 dated

August 9, 1994 describes vagal stimulation to treat sleep disorders. U.S. Pat. No. 5,707,400 dated January 13, 1998 describes vagal stimulation to treat refractory hypertension. U.S. Pat. No. 6,473,644 dated October 29, 2002 describes vagal stimulation to treat heart failure. U.S. Pat. No. 5,571,150 dated November 5, 1996  
5 describes vagal stimulation to treat patients in comas. As previously described, U.S. Pat. No. 5,540,730 dated July 30, 1996 describes vagal stimulation to treat motility disorders and U.S. Pat. No. 6,610,713 dated August 26, 2003 describes vagal stimulation to inhibit inflammatory cytokine production.

All of the foregoing suffer from undesired effects of vagal pacing on  
10 cardiovascular, gastrointestinal or other organs. Nerve conduction blocking permits longer pulse durations which would otherwise have adverse effects on other organs such as those of the cardiovascular or gastrointestinal systems. In accordance with the present invention, all of the foregoing disclosures can be modified by applying a blocking electrode and blocking signal as disclosed herein to prevent adverse side  
15 effects. By way of specific example, pacing a vagus nerve in the thoracic cavity or neck combined with a blocking electrode on the vagus nerve distal to the pacing electrode can be used to treat neuropsychiatric disorders (such as depression and schizophrenia) and Parkinson's and epilepsy and dementia. In such treatments, the blocking electrode is placed distal to the stimulating electrode 25 shown in Figs. 4 and 2, respectively, of each of U.S. Pat. Nos. 5,269,303 and 5,299,569. The present  
20 invention thereby enables the teachings of the afore-referenced patents listed in foregoing two paragraphs.

As described, the parameters of the stimulating and blocking electrodes can be inputted via a controller and, thereby, modified by a physician. Also, Fig. 2  
25 illustrates a feedback for controlling a stimulating electrode. Feedbacks for stimulating electrodes are also described in the patents. The blocking electrode can also be controlled by an implanted controller and feedback system. For example, physiologic parameters (e.g., heart rate, blood pressure, etc.) can be monitored. The blocking signal can be regulated by the controller to maintain measured parameters  
30 in a desired range. For example, blocking can be increased to maintain heart rate within a desired rate range during stimulation pacing.

#### **9. Opportunity for Physician to Alter Treatment for Specific Patient**

35 Gastrointestinal disorders are complex. For many, the precise mechanism is of the disorder is unknown. Diagnosis and treatment are often iterative processes. The present invention is particularly desirable for treating such disorders.

Use of proximal and distal blocking electrodes in combination with one or more pacing electrode permits a physician to alter an operating permutation of the electrodes. This permits regional and local up- or down-regulation of the nervous system and organs. Further, pacing parameters (duty cycle, current, frequency, pulse length) can all be adjusted. Therefore, the treating physician has numerous options to alter a treatment to meet the needs of a specific patient.

In addition, a physician can combine the present invention with other therapies (such as drug therapies like prokinetic agents).

## 10 D. Alternative Embodiments

### 1. Background

With reference to Fig. 9, a stomach S is shown schematically for the purpose of facilitating an understanding of alternative embodiments of the invention as illustrated in Figs. 10-15. In Fig. 9, the stomach S is shown with a collapsed fundus F which is deflated due to fasting. In practice, the fundus F can be reduced in size and volume (as shown in Fig. 9) or expanded (as shown in Fig. 12).

The esophagus E passes through the diaphragm D at an opening or hiatus H. In the region where the esophagus E passes through the diaphragm D, trunks of the vagal nerve (illustrated as the anterior vagus nerve AVN and posterior vagus nerve PVN) are disposed on opposite sides of the esophagus E. It will be appreciated that the precise location of the anterior and posterior vagus nerves AVN, PVN relative to one another and to the esophagus E are subject to a wide degree of variation within a patient population. However, for most patients, the anterior and posterior vagus nerves AVN, PVN are in close proximity to the esophagus E at the hiatus H where the esophagus E passes through the diaphragm D.

The anterior and posterior vagus nerves AVN, PVN divide into a plurality of trunks that innervate the stomach directly and via the enteric nervous system and may include portions of the nerves which may proceed to other organs such as the pancreas, gallbladder and intestines. Commonly, the anterior and posterior vagus nerves AVN, PVN are still in close proximity to the esophagus E and stomach (and not yet extensively branched out) at the region of the junction of the esophagus E and stomach S.

In the region of the hiatus H, there is a transition from esophageal tissue to gastric tissue. This region is referred to as the Z-line (labeled "Z" in the Figures). Above the Z-line, the tissue of the esophagus is thin and fragile. Below the Z-line, the tissue of the esophagus E and stomach S are substantially thickened and more vascular. Within a patient population, the Z-line is in the general region of the lower



esophageal sphincter. This location may be slightly above, slightly below or at the location of the hiatus H.

## 2. Implanted Band Electrode

### 5 a. Description of Device

With reference to Fig. 10, a band 200 is shown placed around the esophagus E below the diaphragm D and overlying the anterior and posterior vagus nerves AVN, PVN at the cardiac notch CN. Alternatively, it can be placed completely around the upper portion of the stomach near its junction of the esophagus.

10 Placement of a band 200 around the esophagus E directly beneath the diaphragm D ensures that the band may be placed around the anterior and posterior vagus nerves AVN, PVN without the need for extensive dissection of the nerves AVN, PVN. In a preferred embodiment, the nerves AVN, PVN are indirectly stimulated by passing electrical signals through the tissue surrounding the nerves.

15 The band 200 may be formed of polyester or the like or any other suitable material which may be sutured in place or otherwise fastened in place surrounding the esophagus E or gastric cardia. Preferably, the band 200 is placed at the junction of the esophagus E and stomach S such that the band may overlie both the esophagus E and stomach S at the cardiac notch CN.

20 The band 200 may have a plurality of electrodes which, in the embodiment of Fig. 10 include an upper electrode array 202 and a lower electrode array 203. In the embodiment of Fig. 11 (in which a band 200 is shown lying flat), the electrode arrays 202, 203 are shown with electrodes placed at an angle relative to the cylindrical axis X-X of the band 200.

25 Placement of the band 200 as described ensures that at least a subset of the electrodes 202, 203 will be in overlying relation to the anterior and posterior vagus nerves AVN, PVN. As a result, energizing the electrodes 202, 203 will result in stimulation of the anterior and posterior vagus nerves AVN, PVN and/or their branches.

30 In therapeutic applications, the upper array 202 of electrodes may be connected to a blocking electrical signal source (with a blocking frequency and other parameters as previously described) and the lower array 203 of electrodes may be connected to a stimulation electrical signal source as previously described. Of course, only a single array of electrodes could be used with all electrodes connected  
35 to either a blocking or a stimulating signal.

The electrical connection of the electrodes 202, 203 to a controller is not shown but may be as previously described by having a leads connecting the electrodes directly to an implantable controller. Alternatively, and as previously

described, electrodes may be connected to an implanted antenna for receiving a signal to energize the electrodes.

The use of an array of electrodes permits the collar 200 to be placed without the need for great accuracy at the time of placement. In the event it is desirable that electrodes not directly overlying a vagus nerve be deactivated, the electrodes could, through operation of a controller, be individually energized to detect a physiological response. The absence of a physiological response (other than possible muscular action of the stomach and esophagus) would indicate the absence of an overlying relation to a vagus nerve. The presence of a physiological response would indicate overlying relation of the tested electrode to a vagus nerve.

By identifying which electrodes create a physiologic response, the remaining electrodes (i.e., those not having a physiological response) could be permanently deactivated. An example of a physiological response would be a cardiovascular response which may be attributed to a signal of about 2-80 hertz and up to 50 milliamps and as more fully described in US Patent 6,532,388 to Hill et al dated March 11, 2003. As a result, a selected one of the AVN or PVN could be energized.

It will be appreciated the foregoing description of identifying electrodes to be deactivated is a non-limiting embodiment. For example, all electrodes could be energized. The therapies as previously described could be employed by using blocking electrodes or stimulation electrodes or both in order to block or energize (or both) the vagus nerve.

Figure 10 also illustrates an alternative embodiment in the form of a band 200' surrounding the body of the stomach S and having arrays 202', 203'. Since the band 200' is more distal to the esophagus E, different and more distal trunks of the vagus nerves would be energized. Also, such a placement would permit the option of covering the anterior vagus nerve while not covering the posterior vagus nerve (or visa versa).

With the embodiment shown in Fig. 10, the benefits of vagal stimulation with resulting enteric rhythm management and the aforementioned benefits of blocking can be achieved without the need for extensive dissection of the vagus nerve. Further, the benefits can be achieved without the need for directly clamping electrodes on a vagus nerve, thereby reducing the possibility of injury to a vagus nerve.

In addition to the benefits of nerve stimulation, the band 200 can also be used to restrict and potentially lengthen the esophagus thereby reducing possibilities for reflux as more fully described in commonly assigned and co-pending U.S. patent application Ser. No. 10/600,080 filed June 20, 2003 and entitled "Gastro-Esophageal Reflux Disease" (GERD) Treatment Method and Apparatus".

### b. Application to Obesity and Satiety

The embodiment of Fig. 10 is particularly suitable for the treatment of obesity. Obesity is of epidemic proportions and is associated with large decreases in life expectancy and early mortality. Peeters, et al., "Obesity in Adulthood and Its  
5 Consequences for Life Expectancy: A Life Table Analysis", Annals of Internal Medicine, Vol. 138, No. 1, pp. 24 – 32 (2003).

In the embodiment of Fig. 10, the upper band 200 is placed around the stomach near the cardiac notch CN. Electrode array 202 may be de-activated (or not present on the band 200). Lower array 203 can be energized with a blocking signal.

10 The prior art suggests stimulating the vagas with a stimulating signal for treating obesity or eating disorders. See, e.g., U.S. Pat. No. 5,188,104 to Wernicke et al., dated February 23, 1993; U.S. Pat. No. 5,263,480 to Wernicke et al., dated November 23, 1993; U.S. Pat. No. 6,587,719 to Barrett et al., dated July 1, 2003 and U.S. Pat. No. 6,609,025 to Barrett et al., dated August 19, 2003. These patents all  
15 describe stimulating, non-blocking signals (e.g., stimulating to a level slightly below a so-called "retching threshold" as described in the '025 patent). As such, all fail to note the problem associated with obesity and eating discords that is not addressed by stimulating the vagus but, rather, by blocking stimulation on the vagus.

The blocking at cardiac notch CN reduces fundal accommodation and creates  
20 satiety sensations. Such a physiologic response is suggested by vagotomy data in which truncal vagotomy patients have experienced weight loss and increased satiety. See, e.g., Kral, "Vagotomy as a Treatment for Morbid Obesity", Surg. Clinics of N. Amer., Vol. 59, No. 6, pp. 1131 – 1138 (1979), Gortz, et al., "Truncal Vagotomy Reduces Food and Liquid Intake in Man", Physiology & Behavior, Vol. 48, pp. 779  
25 – 781 (1990), Smith, et al., "Truncal Vagotomy in Hypothalamic Obesity", The Lancet, pp. 1330 – 1331 (1983) and Kral, "Vagotomy for Treatment of Severe Obesity", The Lancet, pp. 307 – 308 (1978).

The optional lower band 200' is placed lower on the stomach (e.g., close to the pylorus). The lower electrode array 203' of the lower band 200' is energized  
30 with a stimulation signal to modulate intestinal motility in the event motility is otherwise impaired by the upper band blocking.

The upper array 202' of the lower band 200' is energized with a blocking signal so that the stimulation signal at electrodes 203' does not interfere with the blocking effect of electrodes 203 of upper band 200. In this obesity treatment, the  
35 electrodes of the bands 200, 200' can be placed on constricting bands (such as the well-known Lap-Band® system of Inamed Inc., Santa Barbara, CA, USA, and used in obesity treatment). More preferably, the bands 200, 200' are not constricting thereby minimizing erosion risks otherwise associated with highly constricting

bands. However, the neural blocking technology of the present invention can be incorporated into such constricting bands or used in conjunction other obesity surgeries or therapies. Specifically, the scientific literature indicates a vagotomy in combination with other obesity procedure (e.g., antrectomy, gastropasty and biliopancreatic bypass) improves weight loss procedures. Tzu-Ming, et al., "Long-Term Results of Duodenectomy with Highly Selective Vagotomy in the Treatment of complicated Duodenal Ulcers", Amer. J. of Surg., Vol. 181, pp. 372 – 376 (2001), Kral, et al., "Gastropasty for Obesity: Long-Term Weight Loss Improved by Vagotomy", World J. Surg., Vol. 17, pp. 75 – 79 (1993), and Biron, et al., "Clinical Experience with Biliopancreatic Bypass and Gastrectomy or Selective Vagotomy for Morbid Obesity", Canadian J. of Surg., Vol. 29, No. 6, pp. 408 – 410 (1986).

Vagal neural blocking simulates a vagotomy but, unlike a vagotomy, is reversible and controllable. Therefore, while obesity is particularly described as a preferred treatment, the vagal neural block of the present invention can be used as a less drastic procedure for treatments previously performed with a vagotomy. Without limitation, these include obesity, ulcers or chronic pain or discomfort (alone or in combination with conjunctive procedures).

Further, bulimia has been identified as a disease amenable to treatment by decreasing afferent vagal activity via pharmacological vagal inhibitors delivered systemically. Faris, et al., "Effect of Decreasing Afferent Vagal Activity with Ondansetron on Symptoms of Bulimia Nervosa: a Randomized, Double-Blind Trial", The Lancet, pp. 792 – 797 (2000). Therefore, bulimia and other diseases treatable with vagal blocker drugs can be treated with the targeted and site-specific vagal neural block of the present invention.

### 3. Acute Treatment Device

#### a. Device Description

Figure 12 illustrates a still further embodiment of the present invention where a nasogastric tube 300 is passed into the stomach. It will be appreciated that nasogastric tubes are well known and form no part of this invention per se. Some nasogastric tubes have specialized functions. An example is a tamponade tube having gastric and esophageal balloons. An example of such is the Bard® Minnesota Four Lumen Esophagogastric Tamponade Tube for the Control of Bleeding from Esophageal Varices as described in product literature (information for use) contained with the product of that name dated 1998 by C. R. Bard, Inc., Covington, GA, USA. Further, while a nasogastric tube is a preferred embodiment other devices (e.g., an orogastric tube or any elongated device to position electrodes near the esophagus/stomach junction) could be used. Also, while placement at the

esophagus/stomach junction is preferred, the device can be placed in a different lumen (e.g., the trachea) for transmucosal stimulation.

The nasogastric tube 300 is multi-lumen tube which includes distal openings 302 to which suction can be applied to remove gastric contents through the tube 300.  
5 A compliant balloon 304 surrounds the gastric tube. Proximal to the balloon 304 is an opening 309 in communication with a lumen (not shown) to which a suction can be applied to remove saliva through the opening 309.

The balloon 304 has a plurality of electrodes which may include an upper array 306 of electrodes and a lower array 307 of stimulation electrodes. The  
10 electrodes of the upper array 306 may be connected to a blocking signal source via conductors 306a (Fig. 13). The electrodes of the lower array 307 may be connected to a stimulation signal source via conductors 307a. The conductors 306a, 307a may be passed through a lumen in the tube 300 to an external controller (not shown). As a result, multiple electrodes can be energized for transmucosal stimulation of the  
15 anterior and posterior vagus nerves AVN, PVN. Fig. 14 shows an alternative design where the arrays 306, 307 are replaced with expandable, circumferential electrodes 306', 307' connected to a controller (not shown) by conductors 306a', 307a'.

As in the embodiment of Fig. 10, the individual electrodes of the arrays 306, 307 may optionally be selectively energized to detect a cardiovascular signal  
20 indicating an electrical coupling of the electrodes to the vagus nerves AVN, PVN. Electrodes that do not create such a coupling may optionally be deactivated such that only the electrodes having an effective coupling with the vagus nerves AVN, PVN will be activated. Also, and as in the embodiment of Fig. 10, there may be a single array of electrodes or all electrodes may be energized with either a blocking or  
25 stimulation signal.

It will be noted in this embodiment that the electrodes are disposed abutting the mucosal surface of the esophageal and stomach lining and are not in direct contact with the vagus nerves AVN, PVN. Instead, the electrodes are spaced from the vagus nerves AVN, PVN by the thickness of the stomach and lower esophageal  
30 wall thickness.

Transmucosal electrical stimulation of nerves is well known. Such stimulation is disclosed in U.S. Pat. No. 6,532,388 to Hill et al dated March 11, 2003 (describing transmucosal stimulation of nerves across a trachea using a balloon with electrodes in the trachea to modulate cardiac activity). Also, the phenomena of  
35 transmucosal electrical stimulation of nerves is described in Accarino, et al, "Symptomatic Responses To Stimulation Of Sensory Pathways In The Jejunum", Am. J. Physiol., Vol. 263, pp. G673-G677 (1992) (describing afferent pathways inducing perception selectively activated by transmucosal electrical nerve

- stimulation without disruption of intrinsic myoelectrical rhythm); Coffin, et al, "Somatic Stimulation Reduces Perception Of Gut Distention In Humans", Gastroenterology, Vol. 107, pp. 1636-1642 (1994); Accarino, et al, "Selective Dysfunction Of Mechano Sensitive Intestinal Afferents In Irritable Bowel Syndrome", Gastroenterology, Vol. 108, pp. 636-643 (1994), Accarino, et al "Modification Of Small Bowel Mechanosensitivity By Intestinal Fat", GUT, Vol. 48, pp. 690-695 (2001); Accarino, et al, "Gut Perception In Humans Is Modulated By Interacting Gut Stimuli", Am. J. Physiol. Gastrointestinal Liver Physiol., Vol. 282, pp. G220-G225 (2002) and Accarino, et al, "Attention And Distraction Colon Affects On Gut Perception", Gastroenterology, Vol. 113, pp. 415-442 (1997).

Alternative embodiments of the transmucosal stimulation device of Fig. 12 are shown in Fig. 15 and 16. In Fig. 15, the balloon 304' is conical in shape with a base end 304a' placed distally on the tube 300'. After expansion, the base end 304a' expands within the stomach S. The physician then pulls on the tube 300'. The base end 304a' (which is larger in diameter than the esophagus E) abuts the stomach S at the cardiac notch CN acting as a stop. This insures the electrodes 305' (only a single array is shown for ease of illustration) abuts the mucosal tissue at the junction of the stomach S and esophagus E. The electrodes 305' are on the narrow end 304b' of the balloon 304' and expansion of the balloon 304' ensures contact of the electrodes with the mucosal tissue.

Fig. 16 illustrates an embodiment using two balloons 304'' and 309''. The distal balloon 309'', when expanded, is larger than the esophagus E and acts as a stop when the physician pulls on the tube 300''. The electrodes 305'' are on a smaller balloon 304'' which may expand in the esophagus E. The balloon 304'', 309'' are positioned for the electrodes 305'' to be against the mucosal tissue at the junction of the stomach S and esophagus E when the distal balloon 309'' abuts the cardiac notch CN and the proximal balloon 304'' is expanded. The electrodes may be positioned to be completely within the stomach to reduce risk of injury to esophageal tissue. More conveniently, a tube such as the afore-mentioned Bard® tube may be modified for electrodes to be placed on the proximal side of the gastric balloon.

In all of the foregoing, a balloon is expanded to urge the electrodes against the mucosal tissue. While this is a presently preferred embodiment, any mechanism for urging the electrodes against the mucosal tissue may be used. In each of Figs. 15 and 16, the tube 300', 300'' is shown as it passes through the balloons 304', 304'' and 309''. This illustration is made to indicate the tube passes through the balloons and does terminate at the balloons. In fact, as the tube 300', 300'' passes through

the balloons 304', 304'' and 309'' it would be surrounded by the material of the balloons 304', 304'' and 309'' and would not be visible.

A still further embodiment is shown in Fig. 17. Instead of directly stimulating with current, the nerves are stimulated with magnetic fields. In this case, the electrodes are coils 307''' insulated within the balloon 304'''. The coils 307''' create magnetic fields which inductively couple with the vagus nerves to create the blocking and stimulating impulses within the nerves.

#### **b. Application to Acute Pancreatitis**

When energized with a blocking frequency, the embodiment of Fig. 13 is useful for treating acute or recurrent pancreatitis. This extremely serious disease is characterized by an over-active pancreas which excretes digestive enzymes to such an extent that the pancreas itself is digested. The disease can be extremely painful. In many cases, the disease is fatal. The number of US patients who suffer an episode of acute pancreatitis is approximately 185,000 annually. Baron, et al., "Acute Necrotizing Pancreatitis", New England J. of Medicine, Vol. 340, No. 18, pp. 1412 – 1417 (1999). This high incidence, coupled with the cost and length of stay required, make the total cost of this disease to society enormous. No definitive therapy is currently available to treat these patients except supportive care. Furthermore, the overall mortality rate for severe pancreatitis is about 20 to 30%. Id.

A recent study reported that the average total hospital cost to obtain a survivor of severe, acute pancreatitis is nearly \$130,000 with an average length of hospital stay of 40 days. Soran, et al., "Outcome and quality of life of patients with acute pancreatitis requiring intensive care", J. Surg. Res., 91(1), pp. 89 – 94 (2000). Further complicating the management of these patients is the uncertainty surrounding the prognosis because the course of the disease is unpredictable at initial presentation. Chatzicostas, et al., "Balthazar computed tomography severity index is superior to Ranson criteria and APACHE II and II scoring systems in predicting acute pancreatitis outcome", J. Clinical Gastroenterology, 36(3), pp. 253 – 260 (2003). If patients could be successfully treated during the initial phases of the disease, with a higher survival rate, there is a high probability of returning to a productive life. Soran, et al., supra.

Pancreatitis may be associated with a number of etiologies including chronic alcoholism or gallstones (e.g., gallstones lodged in the pancreatic or common duct). When acute pancreatitis becomes severe, treatment options are severely limited. Morbidity and mortality rates for pancreatitis are sobering. Baron, et al., "Acute Necrotizing Pancreatitis", New England J. of Medicine, Vol. 340, No. 18, pp. 1412 –

1417 (1999) and Steer et al., "Chronic Pancreatitis", New England J. of Medicine, pp. 1482 – 1490 (1995).

Down-regulating vagal activity can be used to treat pancreatitis. A recently reported finding in experimental pancreatitis demonstrated that the vagus nerves are strongly implicated in the pathophysiology of pancreatitis. Yoshinaga, et al., "Cholecystokinin Acts as an Essential Factor in the Exacerbation of Pancreatic Bile Duct Ligation-Induced Rat Pancreatitis Model Under Non-Fasting Condition", Japanese J. Pharmacol., Vol. 84, pp. 44 – 50 (2000). Pharmacologic means of decreasing pancreatic secretion have been attempted with limited success because of the dose-limiting side effects encountered with the drugs, their lack of specificity or their lack of availability. In fact, one recent trial of a specific blocker of parasympathetic (vagus nerves) control of secretion demonstrated a shortened recovery period in patients with acute pancreatitis while trials with other pancreatic down-regulating drugs that are less specific or potent have proven to be disappointing. Zapater, et al., "Do Muscarinic Receptors Play a Role in Acute Pancreatitis?", Clin. Drug Invest., 20(6), pp. 401 – 408 (2000); Norton, et al., "Optimizing Outcomes in Acute Pancreatitis", Drugs, 61(11), pp. 1581 – 1591 (2001). Atropine is a drug that blocks parasympathetic nerve endings. It is known to be desirable to use atropine in acute pancreatitis patients to down-regulate pancreatic activity. Unfortunately, for most such patients, this drug cannot be used due to its many side effects.

Acute pancreatitis patients may be placed on intravenous feeding with the device 300 left in place for a chronic length of time (e.g., several days or weeks). At least the electrodes of the lower array 307 may be energized with a blocking signal for the treatment of acute pancreatitis. The invention permits down-regulation of pancreatic output through vagal blocking without the need for undesirable surgery for direct vagal access.

In addition to utility for treating pancreatitis, the present invention may be used to avoid pancreatitis in patients having an increased likelihood of developing the disease. For example, patients undergoing endoscopic retrograde cholangiopancreatography (ERCP) and/or related procedures are known to having a higher likelihood of developing pancreatitis. Such patients may be treated with the present invention with a blocking signal to down-regulate pancreatic output and reduce the likelihood of developing pancreatitis.

Many physicians treating patients with pancreatitis use a nasogastric tube as part of the treatment. As a result, the present invention is illustrated as being incorporated on a nasogastric tube. However, a significant body of physicians adheres to a belief that pancreatitis patients benefit from a feeding involving placing



nourishment directly into the jejunum portion of the small intestine via a naso-jejunal tube. While the present invention is illustrated in an embodiment of placement of the balloon and electrodes on a naso-gastric tube, the invention can also be placed on a nasojejunal tube or a nasogastricjejunal tube.

5

### c. Application to Ileus

With the device of Figs. 12 – 16, the distal electrodes 307 may be energized with a stimulation frequency as described for treatment of ileus. Post-trauma and post-surgery, patients may experience ileus which is a dysfunction of the GI tract characterized in part by a lack of motility through the intestines. Prolonged ileus can result in stasis and serious infection. Kaiser, “Gallstone Ileus”, New England J. of Medicine, Vol. 336, No. 12, pp. 879 – 880 (1997) (correspondence), Taguchi, et al., “Selective Postoperative Inhibition of Gastrointestinal Opioid Receptors”, New England J. of Medicine, Vol. 345, No. 13, pp. 935 – 940 (2001) and Steinbrook, “An Opioid Antagonist For Postoperative Ileus”, New England J. of Medicine, Vol. 345, No. 13, pp. 988 – 989 (2001) (Editorial).

Ileus patients commonly have nasogastric tubes as a regular part of their hospital stay. Without additional invasiveness, the present invention can be used as the nasogastric tube with the addition of stimulation electrodes to stimulate the vagus to enhance motility.

The embodiment of Fig. 12 would permit the lower electrodes 307 to be energized for stimulation frequency to treat ileus. Optionally, the upper electrodes 306 can be energized for blocking frequency if needed to prevent antidromic inhibitory responses or to prevent undesired cardiac response.

The embodiment of Fig. 12 – 16 is useful to permit a diagnosis for a surgical implants as described in foregoing embodiments. Namely, responsiveness of a patient’s gastro-intestinal symptoms (such as IBS) to the embodiment of Fig. 12 could justify a more invasive surgical placement of electrodes directly on the anterior or posterior vagus nerves.

In Fig. 15, a blocking or stimulating signal can be applied to the electrode 305’. A blocking frequency is anticipated to be a therapeutic value for treating, for example, acute pancreatitis or an exacerbation of chronic pancreatitis. A stimulating frequency is anticipated to be of therapeutic value for treating ileus. With the embodiment of Fig. 12, ileus, for example, can be treated by applying a stimulation frequency to the lower electrode 307. A blocking frequency to the upper electrode 306 can be used to block antidromic responses or to block adverse side effects of the stimulation signal on proximal organs (e.g., cardiac responses).

#### 4. Diagnostic Device

Fig. 18 illustrates a still further embodiment of the present invention where a stimulating electrode 400 is placed near a distal end of an esophageal gastric duodenal (EGD) scope 402. Leads (not shown) pass through the scope 402 connecting the electrode 400 to a controller (not shown).

Stimulation may be applied via the electrode 400 for transmucosal stimulation of an opposing vagus nerve (AVN in Fig. 18). Proper placement to achieve stimulation can be identified through the previously described techniques of identifying a cardiovascular response to the stimulation indicating appropriate opposition of the electrode 400 to a vagus nerve.

The scope may be left in place or may be removed by placing the electrode attached to the mucosal wall (Fig. 19) through a pigtail or other attachment (such as an adhesive) with leads passing through the nostril or mouth. Alternatively, the electrode 400 may be kept in place positioned underneath the mucosal layer and may be energized by radio frequencies applied externally.

It will be appreciated that attachment of electrical apparatus to internal mucosal layers of patients' is well known. Such a system is described with respect to the Bravo™ pH monitoring system of Medtronic, Inc., Minneapolis, Minnesota, USA and as described in its product literature UC 200300235 EN N15344 (2002) titled "Bravo™ pH Monitoring System Catheter-Free pH Testing".

The foregoing embodiment is particularly useful for identifying patients responsive to blocking and stimulation as a diagnostic before applying a more invasive procedure using the blocking and stimulation apparatus and methods described herein.

With the foregoing detailed description of the present invention, it has been shown how the objects of the invention have been attained in a preferred manner. Modifications and equivalents of disclosed concepts such as those which might readily occur to one skilled in the art, are intended to be included in the scope of the claims which are appended hereto.

We claim:

1. A method for treating at least one of a plurality of disorders of a patient where the disorders are characterized at least in part by vagal activity innervating at least one of a plurality of alimentary tract organs of said patient at an innervation site, said method comprising:
  - positioning a neurostimulator carrier within a body lumen of said patient and with an electrode disposed on said carrier and positioned at a mucosal layer of said lumen in a region of a vagal trunk;
  - applying an electrical signal to said stimulator to modulate vagal activity by an amount selected to treat said disorder.
2. A method according to claim 1 wherein said electrical signal is applied at a frequency selected for said signal to create a neural conduction block to a vagus nerve of said patient at a blocking site proximal to said innervation site with said neural conduction block selected to at least partially block nerve impulses on said vagus nerve at said blocking site.
3. A method according to claim 2 wherein application of said neural conduction block is variable by a controller to alter a characteristic of said block.
4. A method according to claim 2 wherein said at least one of a plurality of disorders is pancreatitis.
5. A method according to claim 4 wherein said neural conduction block is regulated to reduce a pancreatic output of said patient.
6. A method according to claim 2 wherein said at least one of a plurality of disorders is obesity.
7. A method according to claim 6 wherein said neural conduction block is regulated to heighten a sensation of satiety of said patient.
8. A method according to claim 2 wherein said at least one of a plurality of disorders is constipation.
9. A method according to claim 2 wherein said at least one of a plurality of disorders is irritable bowel syndrome or inflammatory bowel disease.

10. A method according to claim 2 wherein said at least one of a plurality of disorders is vagally mediated pain.
- 5 11. A method according to claim 2 wherein said at least one of a plurality of disorders is bulimia.
12. A method according to claim 2 wherein said at least one of a plurality of disorders is ulcers.
- 10 13. A method according to claim 1 wherein said lumen is a trachea of said patient.
14. A method according to claim 1 wherein said lumen is an esophagus of said patient.
- 15 15. A method according to claim 1 wherein said lumen is an interior of a stomach of said patient.
16. A method according to claim 1 wherein said electrical signal is applied at a frequency selected for said signal to create a stimulation with an up-regulation of a vagus nerve of said patient at a stimulation site proximal to said innervation site.
- 20 17. A method according to claim 16 wherein said at least one of a plurality of disorders is ileus.
- 25 18. An apparatus for treating at least one of a plurality of disorders of a patient where the disorders are characterized at least in part by vagal activity innervating at least one of a plurality of alimentary tract organs of said patient at an innervation site, said apparatus comprising:
- 30 a neurostimulator carrier sized to be passed through a selected body lumen of said patient and having an electrode disposed on said carrier to be positioned at a mucosal layer of said lumen;
- 35 a source of electrical signal electrically coupled to said electrode to modulate vagal activity by an amount selected to treat said disorder.

19. An apparatus according to claim 18 wherein said source is selected to generate a blocking signal selected to at least partially block nerve impulses on said vagus nerve at said blocking site.
- 5 20. An apparatus according to claim 19 comprising a controller for selectively controlling parameters of said blocking signal.
21. An apparatus according to claim 20 wherein said controller is inductively coupled to said electrode to electrically control said electrode remote from an interior of said patient's body.
- 10 22. An apparatus according to claim 18 wherein said carrier is sized to pass into a trachea of said patient.
- 15 23. An apparatus according to claim 18 wherein said carrier is sized to pass into an esophagus of said patient.
24. An apparatus according to claim 18 wherein said carrier is sized to pass into an interior of a stomach of said patient.

20

**FIG. 1**






FIG.2




**Fig. 3**  
**Low Vagal and Enteric Tone Before Pacing**







**Fig. 4**  
Pacing Without Block



**Fig. 5**  
**Implantable System**  
*Beta Medical - Vagus Nerve Stimulation System*





**Fig. 6**  
Pacing With Proximal Block
























FIG. 18



<sup>19</sup>  
FIG. 19

# INTERNATIONAL SEARCH REPORT

Int ional Application No

PC1/US2004/002849

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61N1/36

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2002/087192 A1 (BARRETT BURKE T ET AL) 4 July 2002 (2002-07-04) abstract; figures 4,5 paragraph '0014! - paragraph '0025! paragraphs '0035!, '0042!, '0043! ---	18-24
Y	US 5 292 344 A (DOUGLAS DONALD D) 8 March 1994 (1994-03-08) abstract; figures 2A,2B,2C column 3, line 40 -column 4, line 12 column 5, line 4 - line 43 ---	18-24
Y	US 5 263 480 A (BAKER JR ROSS G ET AL) 23 November 1993 (1993-11-23) abstract; figure 2 column 3, line 55 -column 7, line 25 column 10, line 1 - line 14 column 14, line 41 - line 54 ---	19-21
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the International filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the International filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

22 June 2004

Date of mailing of the international search report

02/07/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Wetzig, T

# INTERNATIONAL SEARCH REPORT

In ... onal Application No  
PC 1/US2004/002849

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 02/26320 A (MEDTRONIC INC) 4 April 2002 (2002-04-04) the whole document -----	22,23

# INTERNATIONAL SEARCH REPORT

national application No.  
PCT/US2004/002849

## Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-17  
because they relate to subject matter not required to be searched by this Authority, namely:  
Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; It is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US2004/002849

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2002087192 A1	04-07-2002	CA 2433376 A1 EP 1357970 A2 WO 02062291 A2 US 2004039427 A1	15-08-2002 05-11-2003 15-08-2002 26-02-2004
US 5292344 A	08-03-1994	AT 191354 T AU 662076 B2 AU 4666993 A CA 2138792 A1 DE 69328304 D1 DE 69328304 T2 EP 0656796 A1 JP 2710864 B2 JP 7508662 T WO 9401172 A1	15-04-2000 17-08-1995 31-01-1994 20-01-1994 11-05-2000 14-12-2000 14-06-1995 10-02-1998 28-09-1995 20-01-1994
US 5263480 A	23-11-1993	US 5188104 A AU 657736 B2 AU 1346992 A CA 2100872 A1 DE 69229499 D1 DE 69229499 T2 EP 0569522 A1 WO 9213592 A1	23-02-1993 23-03-1995 07-09-1992 02-08-1992 05-08-1999 04-05-2000 18-11-1993 20-08-1992
WO 0226320 A	04-04-2002	US 6532388 B1 EP 1322382 A1 WO 0226320 A1 US 2002032468 A1	11-03-2003 02-07-2003 04-04-2002 14-03-2002